# **Introduction - Chemistry**

The following released test questions are taken from the Chemistry Standards Test. This test is one of the California Standards Tests administered as part of the Standardized Testing and Reporting (STAR) Program under policies set by the State Board of Education.

All questions on the California Standards Tests are evaluated by committees of content experts, including teachers and administrators, to ensure their appropriateness for measuring the California academic content standards in Chemistry. In addition to content, all items are reviewed and approved to ensure their adherence to the principles of fairness and to ensure no bias exists with respect to characteristics such as gender, ethnicity, and language.

This document contains released test questions from the California Standards Test forms in 2003, 2004, 2005, and 2006. First on the pages that follow are lists of the standards assessed on the Chemistry Test. Next are released test questions. Following the questions is a table that gives the correct answer for each question, the content standard that each question is measuring, and the year each question last appeared on the test. Reference sheets, provided for students taking the test, are also included as they are necessary in answering some of the questions. It should be noted that asterisked (\*) standards found in the *Science Content Standards for California Public Schools, Kindergarten through Grade 12*, are not assessed on the California Standards Tests in Science and, therefore, are not represented in these released test questions.

The following table lists each reporting cluster, the number of items that appear on the exam, and the number of released test questions that appear in this document. The released test questions for Biology, Chemistry, Earth Science, and Physics are the same test questions found in different combinations on the Integrated Science 1, 2, 3, and 4 tests.

REPORTING CLUSTER	NUMBER OF QUESTIONS ON EXAM	NUMBER OF RELEASED TEST QUESTIONS
Investigation and Experimentation (Standards: CHIE1. a-n)	6	6
Atomic and Molecular Structure  Atomic and Molecular Structure (Standards: CH1. a-e)  Nuclear Processes (Standards: CH11. a-e)	8	8
Chemical Bonds, Biochemistry  Chemical Bonds (Standards: CH2. a-e)  Organic Chemistry and Biochemistry (Standards: CH10. a-c)	) 9	9
Kinetics, Thermodynamics  Gases and Their Properties (Standards: CH4. a-f)  Solutions (Standards: CH6. a-d)  Chemical Thermodynamics (Standards: CH7. a-d)	14	15
Chemical Reactions  Acids and Bases (Standards: CH5. a-d)  Reaction Rates (Standards: CH8. a-c)  Chemical Equilibrium (Standards: CH9. a-b)	13	13
Conservation of Matter and Stoichiometry (Standards: CH3. a-e)	10	9
TOTAL	60	60

In selecting test questions for release, three criteria are used: (1) the questions adequately cover a selection of the academic content standards assessed on the Chemistry Test; (2) the questions demonstrate a range of difficulty; and (3) the questions present a variety of ways standards can be assessed. These released test questions do not reflect all of the ways the standards may be assessed. Released test questions will not appear on future tests.

For more information about the California Standards Tests, visit the California Department of Education's Web site at <a href="http://www.cde.ca.gov/ta/tg/sr/resources.asp">http://www.cde.ca.gov/ta/tg/sr/resources.asp</a>.

#### THE INVESTIGATION AND EXPERIMENTATION REPORTING CLUSTER

The following 14 California content standards are included in the Investigation and Experimentation reporting cluster and are represented in this booklet by six test questions. These questions represent only some ways in which these standards may be assessed on the California Chemistry Standards Test.

Investigation	on and Experimentation
CHIE1.	Scientific progress is made by asking meaningful questions and conducting careful investigations. As a basis for understanding this concept and addressing the content in the other five reporting clusters, students should develop their own questions and perform investigations. Students will:
CHIE1. a.	Select and use appropriate tools and technology (such as computer-linked probes, spreadsheets, and graphing calculators) to perform tests, collect data, analyze relationships, and display data.
CHIE1. b.	Identify and communicate sources of unavoidable experimental error.
CHIE1. c.	Identify possible reasons for inconsistent results, such as sources of error or uncontrolled conditions.
CHIE1. d.	Formulate explanations by using logic and evidence.
CHIE1. e.	Solve scientific problems by using quadratic equations and simple trigonometric, exponential, and logarithmic functions.
CHIE1. f.	Distinguish between hypothesis and theory as scientific terms.
CHIE1. g.	Recognize the usefulness and limitations of models and theories as scientific representations of reality.
CHIE1. h.	Read and interpret topographic and geologic maps.
CHIE1. i.	Analyze the locations, sequences, or time intervals that are characteristic of natural phenomena (e.g., relative ages of rocks, locations of planets over time, and succession of species in an ecosystem).
CHIE1. j.	Recognize the issues of statistical variability and the need for controlled tests.
CHIE1. k.	Recognize the cumulative nature of scientific evidence.
CHIE1. I.	Analyze situations and solve problems that require combining and applying concepts from more than one area of science.
CHIE1. m.	Investigate a science-based societal issue by researching the literature, analyzing data, and communicating the findings. Examples of issues include irradiation of food, cloning of animals by somatic cell nuclear transfer, choice of energy sources, and land and water use decisions in California.
CHIE1. n.	Know that when an observation does not agree with an accepted scientific theory, the observation is sometimes mistaken or fraudulent (e.g., the Piltdown Man fossil or unidentified flying objects) and that the theory is sometimes wrong (e.g., the Ptolemaic model of the movement of the Sun, Moon, and planets).

#### THE ATOMIC AND MOLECULAR STRUCTURE REPORTING CLUSTER

The following 10 California content standards are included in the Atomic and Molecular Structure reporting cluster and are represented in this booklet by eight test questions. These questions represent only some ways in which these standards may be assessed on the California Chemistry Standards Test.

	d Molecular Structure
CH1.	The periodic table displays the elements in increasing atomic number and shows how periodicity of the physical and chemical properties of the elements relates to atomic structure. As a basis for understanding this concept:
CH1. a.	Students know how to relate the position of an element in the periodic table to its atomic number and atomic mass.
CH1. b.	Students know how to use the periodic table to identify metals, semimetals, non-metals, and halogens.
CH1. c.	Students know how to use the periodic table to identify alkali metals, alkaline earth metals and transition metals, trends in ionization energy, electronegativity, and the relative sizes of ions and atoms.
CH1. d.	Students know how to use the periodic table to determine the number of electrons available for bonding.
CH1. e.	Students know the nucleus of the atom is much smaller than the atom yet contains most of its mass.
Nuclear P	
Nucleal F	ocesses
CH11.	Nuclear processes are those in which an atomic nucleus changes, including radioactive decay of naturally occurring and human-made isotopes, nuclear fission, and nuclear fusion. As a basis for understanding this concept:
	Nuclear processes are those in which an atomic nucleus changes, including radioactive decay of naturally occurring and human-made isotopes, nuclear
CH11.	Nuclear processes are those in which an atomic nucleus changes, including radioactive decay of naturally occurring and human-made isotopes, nuclear fission, and nuclear fusion. As a basis for understanding this concept:  Students know protons and neutrons in the nucleus are held together by nuclear forces
CH11. CH11. a.	Nuclear processes are those in which an atomic nucleus changes, including radioactive decay of naturally occurring and human-made isotopes, nuclear fission, and nuclear fusion. As a basis for understanding this concept:  Students know protons and neutrons in the nucleus are held together by nuclear forces that overcome the electromagnetic repulsion between the protons.  Students know the energy release per gram of material is much larger in nuclear fusion or fission reactions than in chemical reactions. The change in mass (calculated by
CH11. a. CH11. b.	Nuclear processes are those in which an atomic nucleus changes, including radioactive decay of naturally occurring and human-made isotopes, nuclear fission, and nuclear fusion. As a basis for understanding this concept:  Students know protons and neutrons in the nucleus are held together by nuclear forces that overcome the electromagnetic repulsion between the protons.  Students know the energy release per gram of material is much larger in nuclear fusion or fission reactions than in chemical reactions. The change in mass (calculated by $E = mc^2$ ) is small but significant in nuclear reactions.  Students know some naturally occurring isotopes of elements are radioactive, as are

#### THE CHEMICAL BONDS, BIOCHEMISTRY REPORTING CLUSTER

The following eight California content standards are included in the Chemical Bonds, Biochemistry reporting cluster and are represented in this booklet by nine test questions. These questions represent only some ways in which these standards may be assessed on the California Chemistry Standards Test.

Chemical	Bonds
CH2.	Biological, chemical, and physical properties of matter result from the ability of atoms to form bonds from electrostatic forces between electrons and protons and between atoms and molecules. As a basis for understanding this concept:
CH2. a.	Students know atoms combine to form molecules by sharing electrons to form covalent or metallic bonds or by exchanging electrons to form ionic bonds.
CH2. b.	Students know chemical bonds between atoms in molecules such as $H_2$ , $CH_4$ , $NH_3$ , $H_2CCH_2$ , $N_2$ , $CI_2$ and many large biological molecules are covalent.
CH2. c.	Students know salt crystals, such as NaCl, are repeating patterns of positive and negative ions held together by electrostatic attraction.
CH2. d.	Students know the atoms and molecules in liquids move in a random pattern relative to one another because the intermolecular forces are too weak to hold the atoms or molecules in a solid form.
CH2. e.	Students know how to draw Lewis dot structures.
Organic C	hemistry and Biochemistry
CH10.	The bonding characteristics of carbon allow the formation of many different organic molecules of varied sizes, shapes, and chemical properties and provide the biochemical basis of life. As a basis for understanding this concept:
CH10. a.	Students know large molecules (polymers), such as proteins, nucleic acids, and starch, are formed by repetitive combinations of simple subunits.
CH10. b.	Students know the bonding characteristics of carbon that result in the formation of a large variety of structures ranging from simple hydrocarbons to complex polymers and biological molecules.
CH10. c.	Students know amino acids are the building blocks of proteins.

#### THE KINETICS, THERMODYNAMICS REPORTING CLUSTER

The following 14 California content standards are included in the Kinetics, Thermodynamics reporting cluster and are represented in this booklet by 15 test questions. These questions represent only some ways in which these standards may be assessed on the California Chemistry Standards Test.

Gases and	d Their Properties
CH4.	The kinetic molecular theory describes the motion of atoms and molecules and explains the properties of gases. As a basis for understanding this concept:
CH4. a.	Students know the random motion of molecules and their collisions with a surface create the observable pressure on that surface.
CH4. b.	Students know the random motion of molecules explains the diffusion of gases.
СН4. с.	Students know how to apply the gas laws to relations between the pressure, temperature, and volume of any amount of an ideal gas or any mixture of ideal gases.
CH4. d.	Students know the values and meanings of standard temperature and pressure (STP).
CH4. e.	Students know how to convert between the Celsius and Kelvin temperature scales.
CH4. f.	Students know there is no temperature lower than 0 Kelvin.
Solutions	
CH6.	Solutions are homogenous mixtures of two or more substances. As a basis for understanding this concept:
CH6. a.	Students know the definitions of solute and solvent.
CH6. b.	Students know how to describe the dissolving process at the molecular level by using the concept of random molecular motion.
CH6. c.	Students know temperature, pressure, and surface area affect the dissolving process.
CH6. d.	Students know how to calculate the concentration of a solute in terms of grams per liter, molarity, parts per million, and percent composition.
Chemical	Thermodynamics
CH7.	Energy is exchanged or transformed in all chemical reactions and physical changes of matter. As a basis for understanding this concept:
CH7. a.	Students know how to describe temperature and heat flow in terms of the motion of molecules (or atoms).
CH7. b.	Students know chemical processes can either release (exothermic) or absorb (endothermic) thermal energy.
СН7. с.	Students know energy is released when a material condenses or freezes and is absorbed when a material evaporates or melts.
CH7. d.	Students know how to solve problems involving heat flow and temperature changes, using known values of specific heat and latent heat of phase change.

#### THE CHEMICAL REACTIONS REPORTING CLUSTER

The following nine California content standards are included in the Chemical Reactions reporting cluster and are represented in this booklet by 13 test questions. These questions represent only some ways in which these standards may be assessed on the California Chemistry Standards Test.

Acids and	Bases					
CH5.	Acids, bases, and salts are three classes of compounds that form ions in water solutions. As a basis for understanding this concept:					
CH5. a.	Students know the observable properties of acids, bases, and salt solutions.					
CH5. b.	Students know acids are hydrogen-ion-donating and bases are hydrogen-ion-accepting substances.					
CH5. c.	Students know strong acids and bases fully dissociate and weak acids and bases partially dissociate.					
CH5. d.	Students know how to use the pH scale to characterize acid and base solutions.					
Reaction	Rates					
CH8.	Chemical reaction rates depend on factors that influence the frequency of collision of reactant molecules. As a basis for understanding this concept:					
CH8. a.	Students know the rate of reaction is the decrease in concentration of reactants or the increase in concentration of products with time.					
CH8. b.	Students know how reaction rates depend on such factors as concentration, temperature, and pressure.					
CH8. c.	Students know the role a catalyst plays in increasing the reaction rate.					
Chemical	Equilibrium					
CH9.	Chemical equilibrium is a dynamic process at the molecular level. As a basis for understanding this concept:					
CH9. a.	Students know how to use LeChatelier's principle to predict the effect of changes in concentration, temperature, and pressure.					
CH9. b.	Students know equilibrium is established when forward and reverse reaction rates are equal.					

#### THE CONSERVATION OF MATTER AND STOICHIOMETRY REPORTING CLUSTER

The following five California content standards are included in the Conservation of Matter and Stoichiometry reporting cluster and are represented in this booklet by nine test questions. These questions represent only some ways in which these standards may be assessed on the California Chemistry Standards Test.

Conservation of Matter and Stoichiometry							
CH3.	The conservation of atoms in chemical reactions leads to the principle of conservation of matter and the ability to calculate the mass of products and reactants. As a basis for understanding this concept:						
CH3. a.	Students know how to describe chemical reactions by writing balanced equations.						
CH3. b.	Students know the quantity one mole is set by defining one mole of carbon 12 atoms to have a mass of exactly 12 grams.						
CH3. c.	Students know one mole equals 6.02 x 10 <sup>23</sup> particles (atoms or molecules).						
CH3. d.	Students know how to determine the molar mass of a molecule from its chemical formula and a table of atomic masses and how to convert the mass of a molecular substance to moles, number of particles, or volume of gas at standard temperature and pressure.						
CH3. e.	Students know how to calculate the masses of reactants and products in a chemical reaction from the mass of one of the reactants or products and the relevant atomic masses.						

- A weather balloon with a 2-meter diameter at ambient temperature holds 525 grams of helium. What type of electronic probe could be used to determine the pressure inside the balloon?
  - A barometric
  - **B** thermometric
  - C calorimetric
  - **D** spectrophotometric

CSC10177

- A scientist observed changes in the gas pressure of one mole of a gas in a sealed chamber with a fixed volume. To identify the source of the changes, the scientist should check for variations in the
  - **A** air pressure outside the chamber.
  - **B** molecular formula of the gas.
  - C temperature of the chamber.
  - **D** isotopes of the gas.

CSC10120

- Blectrical fires cannot be safely put out by dousing them with water. However, fire extinguishers that spray solid carbon dioxide on the fire work very effectively. This method works because carbon dioxide
  - **A** displaces the oxygen.
  - **B** renders the fire's fuel non-flammable.
  - C forms water vapor.
  - **D** blows the fire out with strong wind currents.

CSC00005

- In order to advance to the level of a theory, a hypothesis should be
  - A obviously accepted by most people.
  - **B** a fully functional experiment.
  - C in alignment with past theories.
  - **D** repeatedly confirmed by experimentation.

CSC00144


- Matter is made of atoms that have positive centers of neutrons and protons surrounded by a cloud of negatively charged electrons. This statement is
  - **A** a theory.
  - **B** a hypothesis.
  - C an inference.
  - **D** an observation.

CSC20129

- 6 When a metal is heated in a flame, the flame has a distinctive color. This information was eventually extended to the study of stars because
  - **A** the color spectra of stars indicate which elements are present.
  - **B** a red shift in star color indicates stars are moving away.
  - C star color indicates absolute distance.
  - **D** it allows the observer to determine the size of stars.

## **Chemistry**

#### **Released Test Questions**



Which of the following ordered pairs of elements shows an increase in atomic number but a decrease in average atomic mass?

- A Ag to Pd
- B Co to Ni
- C Ge to Sn
- **D** Cr to Mo

CSC00149

8


Periodic Table of the Elements																		
																	Cl	
						Mn					Ι	Т						
																Te	I	Xe
											Ι							
		1	$\backslash$	Т	Т	Т	Т	Т	Т			Г	Г	Т	Т	$\overline{}$	$\overline{}$	7
			╟	+	+	+	+	+	+	$\dashv$		$\vdash$	$\vdash$	H	+	+	+	┨

Iodine would have chemical properties *most* like

- A manganese (Mn).
- **B** tellurium (Te).
- C chlorine (Cl).
- **D** xenon (Xe).

CSC00028

9



Atomic number increases →

The chart above shows the relationship between the first ionization energy and the increase in atomic number. The letter on the chart for the alkali family of elements is

- A W.
- B X.
- C Y.
- D Z.

CSC00206

- Which of the following atoms has six valence electrons?
  - A magnesium (Mg)
  - **B** silicon (Si)
  - C sulfur (S)
  - **D** argon (Ar)

CSC00185

- Which statement *best* describes the density of an atom's nucleus?
  - A The nucleus occupies most of the atom's volume but contains little of its mass.
  - B The nucleus occupies very little of the atom's volume and contains little of its mass.
  - C The nucleus occupies most of the atom's volume and contains most of its mass.
  - **D** The nucleus occupies very little of the atom's volume but contains most of its mass.



#### Results of Firing Alpha Particles at Gold Foil

Observation:	Proportion:
Alpha particles went straight through gold foil.	> 98%
Alpha particles went through gold foil but were deflected at large angles.	≈ 2%
Alpha particles bounced off gold foil.	≈ 0.01%

# What information do the experimental results above reveal about the nucleus of the gold atom?

- A The nucleus contains less than half the mass of the atom.
- **B** The nucleus is small and is the densest part of the atom.
- C The nucleus contains small positive and negative particles.
- **D** The nucleus is large and occupies most of the atom's space.

CSC20056

- Why are enormous amounts of energy required to separate a nucleus into its component protons and neutrons even though the protons in the nucleus repel each other?
  - A The force of the protons repelling each other is small compared to the attraction of the neutrons to each other.
  - **B** The electrostatic forces acting between other atoms lowers the force of repulsion of the protons.
  - C The interactions between neutrons and electrons neutralize the repulsive forces between the protons.
  - **D** The forces holding the nucleus together are much stronger than the repulsion between the protons.

- A 2-cm-thick piece of cardboard placed over a radiation source would be *most* effective in protecting against which type of radiation?
  - A alpha
  - B beta
  - C gamma
  - D x-ray

CSC00299

- Which of the following is a monatomic gas at STP?
  - A chlorine
  - **B** fluorine
  - C helium
  - D nitrogen

CSC10387

- When cations and anions join, they form what kind of chemical bond?
  - A ionic
  - B hydrogen
  - C metallic
  - D covalent

- Some of the molecules found in the human body are NH<sub>2</sub>CH<sub>2</sub>COOH (glycine), C<sub>6</sub>H<sub>12</sub>O<sub>6</sub> (glucose), and CH<sub>3</sub>(CH<sub>2</sub>)<sub>16</sub>COOH (stearic acid). The bonds they form are
  - A nuclear.
  - **B** metallic.
  - C ionic.
  - D covalent.

CSC10230

- The reason salt crystals, such as KCl, hold together so well is because the cations are strongly attracted to
  - A neighboring cations.
  - **B** the protons in the neighboring nucleus.
  - **C** free electrons in the crystals.
  - **D** neighboring anions.

CSC00150

- Under the same conditions of pressure and temperature, a liquid differs from a gas because the molecules of the liquid
  - A have no regular arrangement.
  - **B** are in constant motion.
  - C have stronger forces of attraction between them.
  - **D** take the shape of the container they are in.

CSC10388

20																
		Р	eri	oc	lic	Ta	abl	e (	of	the	e E	Ξle	m	en	ts	
Г		1														
Г	П	1										Al	П	П		
	Т											Ga	Ge	As		
	T				П			П						П		
	Т	П	Π	Г	Г	Г		П	Г	Г				П		
	Τ	П	Г		Г	Г		П	Г	Г		_				
_		↤	abla	_		_	_		_							

Which of the following elements has the same Lewis dot structure as silicon?

- A germanium (Ge)
- **B** aluminum (Al)
- C arsenic (As)
- D gallium (Ga)

CSC0014

- Which substance is made up of many monomers joined together in long chains?
  - A salt
  - B protein
  - C ethanol
  - **D** propane

CSC00323

- For the polymer, polyvinyl chloride (PVC), ~ CH<sub>2</sub>CH(Cl)CH<sub>2</sub>CH(Cl)CH<sub>2</sub>CH(Cl) ~ the repeating subunit is
  - A CH(Cl).
  - B CH(Cl)CHCH<sub>2</sub>.
  - $\mathbf{C}$   $\mathbf{CH}_{2}\mathbf{CH}$ .
  - $\mathbf{D}$  CH<sub>2</sub>CH(Cl).

## **Released Test Questions**

# Chemistry

- Proteins are large macromolecules composed of thousands of subunits. The structure of the protein depends on the sequence of
  - A lipids.
  - **B** monosaccharides.
  - C amino acids.
  - D nucleosides.

CSC00062

- When someone standing at one end of a large room opens a bottle of vinegar, it may take several minutes for a person at the other end to smell it. Gas molecules at room temperature move at very high velocities, so what is responsible for the delay in detection of the vinegar?
  - **A** the increase in the airspace occupied by vinegar molecules
  - **B** the chemical reaction with nerves, which is slower than other sensory processes
  - C attractive forces between the air and vinegar molecules
  - **D** random collisions between the air and vinegar molecules

CSC00125

- Methane (CH<sub>4</sub>) gas diffuses through air because the molecules are
  - **A** moving randomly.
  - **B** dissolving quickly.
  - C traveling slowly.
  - **D** expanding steadily.

CSC20840

- The volume of 400 mL of chlorine gas at 400 mm Hg is decreased to 200 mL at constant temperature. What is the new gas pressure?
  - **A** 400 mm Hg
  - **B** 300 mm Hg
  - C 800 mm Hg
  - **D** 650 mm Hg

CSC00239

- Standard temperature and pressure (STP) are defined as
  - **A** 0 °C and 1.0 atm pressure.
  - **B** 0 °C and 273 mm Hg pressure.
  - C 0 K and 1.0 atm pressure.
  - D 0 K and 760 mm Hg pressure.

CSC00285

- Under which of the following sets of conditions will a 0.50 mole sample of helium occupy a volume of 11.2 liters?
  - **A** 298 K and 0.90 atm
  - **B** 273 K and 1.10 atm
  - C 373 K and 0.50 atm
  - **D** 273 K and 1.00 atm

CSC10234

- What is the equivalent of 423 kelvin in degrees Celsius?
  - **A** −223 °C
  - **B** −23 °C
  - **C** 150 °C
  - **D** 696 °C

30

SOLUBILITY OF SUBSTANCES IN WATER @ 20 °C										
Substance Formula/State Solubility (g/100g H2O)										
Magnesium chloride	MgCl <sub>2</sub> / solid	54.6								
Ammonia	NH₃ / gas	34.0								
Ethanol	CH <sub>3</sub> CH <sub>2</sub> OH / liquid	infinite								
Benzoic Acid	C <sub>6</sub> H <sub>5</sub> COOH / solid	0.29								

Which of the substances in the table can act as either the solute or the solvent when mixed with 100 grams of water at 20 °C?

- A NH<sub>3</sub>
- B C<sub>6</sub>H<sub>5</sub>COOH
- C MgCl<sub>2</sub>
- D CH<sub>3</sub>CH<sub>2</sub>OH

CSC10055

- If the attractive forces among solid particles are less than the attractive forces between the solid and a liquid, the solid will
  - A probably form a new precipitate as its crystal lattice is broken and re-formed.
  - **B** be unaffected because attractive forces within the crystal lattice are too strong for the dissolution to occur.
  - C begin the process of melting to form a liquid.
  - **D** dissolve as particles are pulled away from the crystal lattice by the liquid molecules.

CSC00088

Water is a polar solvent, while hexane is a nonpolar solvent.

Solute	Water	Hexane
NH <sub>4</sub> CI, ammonium chloride	Soluble	Insoluble
C <sub>10</sub> H <sub>8</sub> , naphthalene	Insoluble	Soluble
C <sub>2</sub> H <sub>5</sub> OH, ethanol	Soluble	Soluble
CO(NH <sub>2</sub> ) <sub>2</sub> , urea	Soluble	Insoluble

Which of the examples above illustrates a nonpolar solute in a polar solvent?

- A NH<sub>4</sub>Cl in water
- $\mathbf{B} \quad \mathbf{C}_{10}\mathbf{H}_8$  in water
- $\mathbf{C}$   $\mathbf{C}_2\mathbf{H}_5\mathbf{OH}$  in hexane
- **D**  $CO(NH_2)_2$  in hexane

CSC20958

- If the solubility of NaCl at 25 °C is 36.2 g/100 g H<sub>2</sub>O, what mass of NaCl can be dissolved in 50.0 g of H<sub>2</sub>O?
  - **A** 18.1 g
  - **B** 36.2 g
  - C 72.4 g
  - **D** 86.2 g

CSC00275

- How many moles of HNO<sub>3</sub> are needed to prepare 5.0 liters of a 2.0 M solution of HNO<sub>3</sub>?
  - **A** 2.5
  - **B** 5
  - **C** 10
  - **D** 20

### **Released Test Questions**

# Chemistry

- The Dead Sea is the saltiest sea in the world. It contains 332 grams of salt per 1000 grams of water. What is the concentration in parts per million (ppm)?
  - A 0.332 ppm
  - **B** 332 ppm
  - C 33,200 ppm
  - **D** 332,000 ppm

CSC20046

- The random molecular motion of a substance is greatest when the substance is
  - A condensed.
  - B a liquid.
  - C frozen.
  - D a gas.

CSC00258

- The boiling point of liquid nitrogen is 77 kelvin. It is observed that ice forms at the opening of a container of liquid nitrogen. The *best* explanation for this observation is
  - A water at zero degrees Celsius is colder than liquid nitrogen and freezes.
  - **B** the nitrogen boils and then cools to form a solid at the opening of the container.
  - C water trapped in the liquid nitrogen escapes and freezes.
  - **D** the water vapor in the air over the opening of the liquid nitrogen freezes out.

CSC00171

- The specific heat of copper is about 0.4 joules/ gram °C. How much heat is needed to change the temperature of a 30-gram sample of copper from 20.0 °C to 60.0 °C?
  - **A** 1000 J
  - **B** 720 J
  - **C** 480 J
  - **D** 240 J


CSC00045

- Equal volumes of 1 molar hydrochloric acid (HCl) and 1 molar sodium hydroxide base (NaOH) are mixed. After mixing, the solution will be
  - A strongly acidic.
  - **B** weakly acidic.
  - C nearly neutral.
  - **D** weakly basic.

## Chemistry

#### **Released Test Questions**





The above picture shows a light bulb connected to a battery with the circuit interrupted by a solution. When dissolved in the water to form a 1.0 molar solution, all of the following substances will complete a circuit allowing the bulb to light *except* 

- A hydrochloric acid.
- **B** sodium nitrate.
- C sucrose.
- **D** ammonium sulfate.

CSC00146

# Which of the following is an observable property of many acids?

- **A** They become slippery when reacting with water.
- **B** They react with metals to release hydrogen gas.
- C They produce salts when mixed with other acids.
- **D** They become more acidic when mixed with a base.

CSC20338

# Potassium hydroxide (KOH) is a strong base because it

- A easily releases hydroxide ions.
- **B** does not dissolve in water.
- **C** reacts to form salt crystals in water.
- **D** does not conduct an electric current.

CSC20341

# Of four different laboratory solutions, the solution with the *highest* acidity has a pH of


- **A** 11.
- **B** 7.
- **C** 5.
- **D** 3.


CSC0017


# A catalyst can speed up the rate of a given chemical reaction by


- **A** increasing the equilibrium constant in favor of products.
- **B** lowering the activation energy required for the reaction to occur.
- C raising the temperature at which the reaction occurs.
- **D** increasing the pressure of reactants, thus favoring products.

Which reaction diagram shows the effect of using the appropriate catalyst in a chemical reaction?









CSC20412

- H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide, naturally breaks down into H<sub>2</sub>O and O<sub>2</sub> over time. MnO<sub>2</sub>, manganese dioxide, can be used to lower the energy of activation needed for this reaction to take place and, thus, increase the rate of reaction. What type of substance is MnO<sub>2</sub>?
  - A a catalyst
  - B an enhancer
  - C an inhibitor
  - D a reactant

CSC1036

- When a reaction is at equilibrium and more reactant is added, which of the following changes is the immediate result?
  - **A** The reverse reaction rate remains the same.
  - **B** The forward reaction rate increases.
  - **C** The reverse reaction rate decreases.
  - **D** The forward reaction rate remains the same.

# Chemistry

## **Released Test Questions**

In which of the following reactions involving gases would the forward reaction be favored by an increase in pressure?

$$\mathbf{A} \quad \mathbf{A} + \mathbf{B} \rightleftharpoons \mathbf{A} \mathbf{B}$$

$$\mathbf{B} \quad \mathbf{A} + \mathbf{B} \rightleftharpoons \mathbf{C} + \mathbf{D}$$

$$\mathbf{C}$$
 2A + B  $\rightleftharpoons$  C + 2D

$$\mathbf{D}$$
 AC  $\rightleftharpoons$  A + C

CSC00129



$$4\mathsf{HCl}_{(g)} + \mathsf{O}_{2(g)} \rightleftarrows 2\mathsf{H}_2\mathsf{O}_{(\mathsf{I})} + 2\mathsf{Cl}_{2(g)} + 113 \; \mathsf{kJ}$$

Which action will drive the reaction to the right?

- A heating the equilibrium mixture
- **B** adding water to the system
- C decreasing the oxygen concentration
- **D** increasing the system's pressure

CSC10082

$$NO_2(g) + CO(g) \rightleftharpoons NO(g) + CO_2(g)$$

The reaction shown above occurs inside a closed flask. What action will shift the reaction to the left?

- A pumping CO gas into the closed flask
- **B** raising the total pressure inside the flask
- C increasing the NO concentration in the flask
- **D** venting some CO<sub>2</sub> gas from the flask

CSC20419

- In a sealed bottle that is half full of water, equilibrium will be attained when water molecules
  - **A** cease to evaporate.
  - **B** begin to condense.
  - C are equal in number for both the liquid and the gas phase.
  - **D** evaporate and condense at equal rates.

52

$$C_3H_8 + O_2 \longrightarrow CO_2 + H_2O$$

This chemical equation represents the combustion of propane. When correctly balanced, the coefficient for water is

- **A** 2.
- **B** 4.
- **C** 8.
- **D** 16.

CSC00311

Which of the following is a balanced equation for the combustion of ethanol (CH<sub>3</sub>CH<sub>2</sub>OH)?

- A  $CH_3CH_2OH + 3O_2 \longrightarrow CO_2 + 2H_2O$
- B  $CH_3CH_2OH + 3O_2 \longrightarrow 2CO_2 + 3H_2O$
- $C \quad CH_3CH_2OH + O_2 \longrightarrow 2CO_2 + 3HO$
- $D \quad CH_3CH_2OH + 2O_2 \longrightarrow 3CO_2 + 2H_2O$

CSC10401

How many moles of carbon-12 are contained in exactly 6 grams of carbon-12?

- **A** 0.5 mole
- B 2.0 moles
- C  $3.01 \times 10^{23}$  moles
- **D**  $6.02 \times 10^{23}$  moles

CSC00068

How many atoms are contained in 97.6 g of platinum (Pt)?

- A  $5.16 \times 10^{30}$
- **B**  $3.01 \times 10^{23}$
- C  $1.20 \times 10^{24}$
- **D**  $1.10 \times 10^{28}$

CSC00255

When methane (CH<sub>4</sub>) gas is burned in the presence of oxygen, the following chemical reaction occurs.

$$CH_4 + 2O_2 \longrightarrow CO_2 + 2H_2O$$

If 1 mole of methane reacts with 2 moles of oxygen, then

- A  $6.02 \times 10^{23}$  molecules of  $CO_2$  and  $6.02 \times 10^{23}$  molecules of  $H_2O$  are produced.
- **B**  $1.2 \times 10^{24}$  molecules of  $CO_2$  and  $1.2 \times 10^{24}$  molecules of  $H_2O$  are produced.
- C  $6.02 \times 10^{23}$  molecules of CO<sub>2</sub> and  $1.2 \times 10^{24}$  molecules of H<sub>2</sub>O are produced.
- **D**  $1.2 \times 10^{24}$  molecules of  $CO_2$  and  $6.02 \times 10^{23}$  molecules of  $H_2O$  are produced.

CSC20428

How many moles of CH<sub>4</sub> are contained in 96.0 grams of CH<sub>4</sub>?

- **A** 3.00 moles
- **B** 6.00 moles
- **C** 12.0 moles
- **D** 16.0 moles

# Chemistry

## **Released Test Questions**

How many atoms are in a chromium sample with a mass of 13 grams?

A  $1.5 \times 10^{23}$ 

**B**  $3.3 \times 10^{23}$ 

C  $1.9 \times 10^{26}$ 

**D**  $2.4 \times 10^{24}$ 

CSC10251



$$Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$$

In this reaction, how many grams of  $Fe_2O_3$  are required to completely react with 84 grams of CO?

**A** 64 g

**B** 80 g

**C** 160 g

**D** 1400 g

CSC00159

$$Mg_3N_2(s) + 6H_2O(l) \rightarrow$$

$$2NH_3(aq) + 3Mg(OH)_2(s)$$

If 54.0 grams of water are mixed with excess magnesium nitride, then how many grams of ammonia are produced?

**A** 1.00

**B** 17.0

C 51.0

**D** 153

Question Number	Correct Answer	Standard	Year of Test
1	A	CHIE1.A	2005
2	С	CHIE1.C	2006
3	A	CHIE1.D	2004
4	D	CHIE1.F	2004
5	A	CHIE1.F	2006
6	A	CHIE1.K	2003
7	В	CH1.A	2004
8	С	CH1.B	2004
9	A	CH1.C	2003
10	С	CH1.D	2003
11	D	CH1.E	2004
12	В	CH1.E	2006
13	D	CH11.A	2005
14	A	CH11.E	2003
15	С	CH2.A	2005
16	A	CH2.A	2006
17	D	CH2.B	2005
18	D	CH2.C	2004
19	С	CH2.D	2005
20	A	CH2.E	2003
21	В	CH10.A	2003
22	D	CH10.A	2006
23	С	CH10.C	2004
24	D	CH4.B	2004
25	A	CH4.B	2006
26	С	CH4.C	2003
27	A	CH4.D	2004
28	D	CH4.D	2006
29	С	CH4.E	2003
30	D	CH6.A	2005
31	D	CH6.B	2004
32	В	CH6.B	2006
33	A	CH6.D	2003
34	С	CH6.D	2004
35	D	CH6.D	2006
36	D	CH7.A	2003
37	D	CH7.C	2004
38	С	CH7.D	2003

Question Number	Correct Answer	Standard	Year of Test
39	C	CH5.A	2003
40	C	CH5.A	2005
41	В	CH5.A	2006
42	A	CH5.C	2005
43	D	CH5.D	2005
44	В	CH8.C	2003
45	D	CH8.C	2005
46	A	CH8.C	2006
47	В	CH9.A	2003
48	A	CH9.A	2004 2005 2006 2005 2004
49	D	CH9.A	
50	C	CH9.A	
51	D	CH9.B	
52	В	CH3.A	
53	В	CH3.A	2005
54	A	CH3.B	2004
55	В	CH3.C	2005
56	C	CH3.C	2006
57	В	CH3.D	2003
58	A	CH3.D	2006
59	C	CH3.E	2005
60	В	СН3.Е	2006

Г			I	<u> </u>	Ι	Γ		E
18 8A	Helium 4.00	10 <b>Ne</b> Neon 20.18	18 <b>Ar</b> Argon 39.95	36 Krypton 83.80	54 <b>Xe</b> Xenon 131.29	86 <b>Bn</b> Radon (222)		71
	17 7A	9 <b>F</b> Fluorine 19.00	17 Chlorine 35.45	35 <b>Br</b> Bromine 79.90	53 — lodine 126.90	85 <b>At</b> Astatine (210)		70 Yb Yterbium 173.04 102 No Nobelium L (259)
	16 6A	8 <b>O</b> Oxygen 16.00	16 Sulfur 32.07	(O)	52 <b>Te</b> Tellurium 127.60	84 <b>Po</b> Polonium (209)		69 <b>Tm</b> Thuliun 168.93 101 <b>Md</b> Mendelevix
	15 5A	7 <b>N</b> itrogen 14.01	15 <b>P</b> Phosphorus 30.97	33 <b>AS</b> Arsenic 74.92	51 <b>Sb</b> Antimony 121.76	83 <b>Bi</b> Bismuth 208.98		68 <b>Er</b> bium 167.26 100 <b>Fm</b> Femium (257)
	14 4A	6 Carbon 12.01	14 Silicon 28.09	Ε	50 <b>Sn</b> Tin 118.71	82 <b>Pb</b> Lead 207.2		67 Holmium 164.93 99 ES Einsteinium (252)
	13 3A	5 <b>B</b> Boron 10.81	13 <b>AI</b> Aluminum 26.98		49 <b>Ln</b> Indium 114.82	81 <b>T</b> Thallium 204.38		66 67  Dy Ho  Dysprosium Holmium 162.50 164.93 98 99  Cf ES  Californium Einsteinium (251) (252)
			12 28	30 <b>Xn</b> Zinc 65.39	48 <b>Cd</b> Cadmium 112.41	80 <b>Hg</b> Mercury 200.59		65 <b>Tb</b> Terbium 158.93 97 97 <b>Bk</b> Berkelium (247)
		Atomic number Element symbol Element name	<del>=</del> = = = = = = = = = = = = = = = = = =	29 <b>Cu</b> Copper 63.55	47 <b>Ag</b> Silver 107.87	79 <b>Au</b> Gold 196.97		64 <b>Gd</b> Gadolinium 157.25 96 Cm Curium (247)
			10	28 Nickel 58.69	46 <b>Pd</b> Palladium 106.42	78 <b>Pt</b> Platinum 195.08		63 Europium 151.96 95 Am Americium (243)
			ic mass* 9 88	27 <b>Co</b> Cobalt 58.93	45 <b>Rh</b> Rhodium 102.91	77 <b>       </b>	109 Mt Meitnerium (268)	Sm Sm Samarium 150.36 94 Pu M Plutonium /
	Key		nic numbe nent symk nent name rage atom 8	26 <b>Te</b> Iron 55.85	44 <b>Bu</b> Ruthenium 101.07	76 <b>Os</b> Osmium 190.23	108 <b>HS</b> Hassium (269)	61 <b>Pm</b> Promethium (145) 93 <b>Np</b> Neptunium (237)
	<b>Ž</b> þ	4++	¬	25 Wn Manganese 54.94	43 <b>Tc</b> Technetium (98)	75 <b>Re</b> Rhenium 186.21	107 <b>Bh</b> Bohrium (264)	60 Nd Neodymium 144.24 92 U
		Na Sodium 22.99	6 6 6 6 8	E C	Molybdenum 95.94	74 <b>W</b> Tungsten 183.84	106 <b>Sg</b> Seaborgium (266)	59   60   61
			5 2 8	23 <b>V</b> Vanadium 50.94	41 <b>Nb</b> Niobium 92.91	73 <b>Ta</b> Tantalum 180.95	105 <b>Db</b> Dubnium (262)	58 Cerium 140.12 90 <b>Th</b> Thorium 232.04
		4 4 84	22 <b>Ti</b> Titanium 47.87	40 <b>Zr</b> Zirconium 91.22	72 <b>Hf</b> Hafnium 178.49	104 <b>Rf</b> Rutherfordium (261)	ue	
		(	8 8 8 8	Scandium 44.96	39 <b>Y</b> ttrium 88.91	57 <b>La</b> Lanthanum 138.91	89 <b>Ac</b> Actinium (227)	antheses, th
	2 2A	<b>Be</b> Beryllium 9.01	12 <b>Mg</b> Magnesium 24.31	20 <b>Ca</b> lcium 40.08	38 Strontium 87.62	56 <b>Ba</b> Barium 137.33	88 <b>Ra</b> Radium (226)	If this number is in parentheses, then it refers to the atomic mass of the most stable isotope.
- ₹	T Hydrogen 1.01	3 <b>Lith</b> ium 6.94	11 Na Sodium 22.99	19 <b>K</b> Potassium 39.10	37 <b>Rb</b> Rubidium 85.47	55 <b>Cs</b> Cesium 132.91	87 <b>Fr</b> Francium (223)	If this number is in p it refers to the atomi most stable isotope.
	-	N	ო	4	Ŋ	9	2	*

Copyright © 2007 California Department of Education

# Chemistry Reference Sheet

Formulas, Constants, and Unit Conversions

**Formulas** 

Calorimetric Formulas –

No Phase Change:  $Q = m(\Delta T)C_p$ 

Combined Gas Law:  $\frac{P_IV_I}{T_I} = \frac{P_2V_2}{T_2}$ 

Ideal Gas Law: PV = nRT

Pressure Formula:  $P = \frac{F}{A}$ 

Mass-Energy Formula:  $E = mc^2$ 

Latent Heat of Fusion:  $Q = m\Delta H_{\mathrm{fus}}$ 

Latent Heat of Vaporization:  $Q = m\Delta H_{\mathrm{vap}}$ 

Constants

Volume of Ideal Gas at STP:  $22.4 \, \frac{\mathrm{L}}{\mathrm{mol}}$ 

Speed of Light in a Vacuum:  $c = 3.00 \times 10^8 \; \frac{\mathrm{m}}{\mathrm{S}}$ 

Specific Heat of Water:  $C_p(\mathrm{H_2O}) = 1.00 \, \frac{\mathrm{cal}}{(\mathrm{g} \, ^{\mathrm{o}}\mathrm{C})} = 4.18 \, \frac{\mathrm{J}}{(\mathrm{g} \, ^{\mathrm{o}}\mathrm{C})}$ 

Latent Heat of Fusion of Water:  $\Delta H_{\rm fus}({\rm H_2O}) = 80~{{\rm cal}\over {\rm g}} = 334~{{\rm J}\over {\rm g}}$ 

Latent Heat of Vaporization of Water:  $\Delta H_{\rm vap}({\rm H_2O}) = 540~{{\rm cal}\over {\rm g}} = 2260~{{\rm J}\over {\rm g}}$ 

**Unit Conversions** 

Calorie-Joule Conversion: 1 cal = 4.184 J

**Absolute Temperature Conversion:**  $K = {}^{\circ}C + 273$ 

**Pressure Conversions:** 1 atm = 760 mm Hg = 760 Torr =  $101.325 \text{ kPa} = 14.7 \frac{1\text{bs.}}{\text{in.}^2} = 29.92 \text{ in. Hg}$